

# Conservation Status of Ornithoptera Croesus, Endemic Butterfly of Bacan Island

*by* Abdu Mas'ud

---

**Submission date:** 04-Oct-2020 12:03PM (UTC+0700)

**Submission ID:** 1404509584

**File name:** s\_of\_Ornithoptera\_Croesus,\_Endemic\_Butterfly\_of\_Bacan\_Island.pdf (419.16K)

**Word count:** 4809

**Character count:** 26535

**4 CONSERVATION STATUS OF *ORNITHOPTERA CROESUS*,  
ENDEMIC BUTTERFLY OF BACAN ISLAND, NORTH MALUKU,  
INDONESIA**

Abdu MAS'UD<sup>1\*</sup>, SUNDARI<sup>1</sup>, Mohamad AMIN<sup>2</sup>,  
Fatchur ROHMAN<sup>2</sup>, Didik WAHYUDI<sup>3</sup>, ALISI<sup>4</sup>

<sup>1</sup>Departement Biology Education – Faculty of Teacher Training and Education – Universitas Khairun – Street Pertamina Campus II Unkhair Gambesi City of South Ternate – North Maluku 97728 Indonesia

<sup>2</sup>Department of Biology – Faculty of Mathematics and Natural Sciences – Universitas Negeri Malang – Street Semarang 5 Sumbersari Lowokwaru District – Malang City East Java 65145 Indonesia

<sup>3</sup>Faculty of Science and Technology – State Islamic University of Maulana Malik Ibrahim – Malang 65145 Indonesia – Street Gajayana Number 50 Dinoyo Lowokwaru District – Malang City East Java 65145 Indonesia

<sup>4</sup>Conservationists Endemic Butterfly Island Bacan North Maluku – Street Oesmansyah Labuha Bacan District South Halmahera Regency North Maluku. 97791, Indonesia

**Abstract**

*Ornithoptera croesus* is a butterfly species that is native and endemic to Bacan Island. This study aimed to update the conservation status of *O. croesus* in Bacan Island based on the current information on the existence, characters, habitat carrying capacity, and conservation status of the species. In this descriptive exploratory study, data were collected through survey, specimen identification, and literature review on the butterfly conservation status. The results of the analyses show that based on the government regulation (PP) No.7/1999, the conservation status of *O. croesus* is protected, while according to IUCN Red List of Threatened Species, this animal is vulnerable, and according to CITES, the status is endangered (Appendix II). This is consistent to the forest conversion and habitat fragmentation phenomena that occur in Mt. Sibela nature reserve area, *O. croesus* natural habitat. Given this condition, the existence of the endemic butterfly has a low species density with adequate environmental capacity in terms of food availability. Both in situ and ex situ conservation efforts should be made to increase *O. croesus* viability

**Keywords:** *O. Croesus*; Vulnerable; IUCN; Endangered; CITES; Appendix II; Endemic

**Introduction**

*Ornithoptera croesus* is one of the endemic butterflies in the Bacan Island. This butterfly can be found in the nature reserve area of the Sibela mountain (CAGS). *O. croesus* is one member of the family Papilionidae. *O. croesus* discovered by Alfred Russel Wallace in 1859 in the expedition Malay Archipelago. This butterfly is named Wallace's Golden Birdwing. *O. croesus* including macrolepidoptera butterfly with the typical body color patterns. *O. croesus* front wing of black males have golden, orange or red front part and with black veins at the edge of the golden yellow wings. On the underside of the wing iridescent green, with grooved black edges. *O. croesus* females have larger wings with dark brown on both upper side and bottom with yellowish and gray spots, black head and chest, with yellow abdomen [1]

\* Corresponding author: abdu.masud@unkhair.ac.id

The existence of the butterfly *O. croesus* in Bacan Island has been known by the local community around [2]. But so far there has not yet been any conservation efforts done both by the community and regional government. Conservation of the endemic butterflies are highly valued compared to that of common butterflies [3]. Endemic species have limited geographical distribution, is very sensitive to the temperature and humidity changes of the forest caused by the forest disruption, thus the local extinction becomes inevitable [4-8]. Other factor that led to the species become endemic is the sensitivity of environmental changes, such as limited tolerance on temperature, humidity and light intensity [9-14]. Some research indicates that the endemic species butterflies have little preference toward human modified habitats [15-17]. In Northern Vietnam, noted that the endemic species of butterflies prefer the nearby forest habitat, while generalist species prefer open habitats [4, 12].

*O. croesus* like other animals also face the threat of scarcity and the extinction, mainly due to conversion of forest lands. There are two types of forest in mountain Sibela conservation namely forest production and limited production forest [18]. Both types of forest are located along the edges of forests in Sibela mountain conservation. Local people have converted the production forest into core estate and around the small holders [18]. The existence of these forests has been eradicated and converted into residential area and community plantation. Most species of butterflies, especially the endemic butterflies (*O. croesus*) rely heavily on one or two types of host plants, so the threat to the host plant jeopardises the existence of the butterflies as well. The decreases and changes in forest ecosystems that occur due to rapid exploitation is a threat to the existence of butterflies in the Bacan Island. For example, the forest area that has abundant of butterflies' life is destroyed and converted into plantations. Although some butterfly population manages to migrate to new habitats, but the availability of food resources for the larvae to grow have vanished, it may include the typical food for butterfly larvae. *O. croesus* requires a stable habitat components, such as the availability of vegetation components as food resources, if there is no or little food resources than their needs, there will be a migration of *O. croesus* to seek new territorial boasts abundance of feed and refuge from the predators.

The main threat of *O. croesus* is the destruction and changes of forest types as the habitat of *O. croesus*. So far, local communities who live along the mount Sibela natural conservation only utilize the potential of forests by exploiting the types of flora resources to meet daily needs, such as rattan, indigo sap, wood, and other forest products. Animals hunting is also carried out, for example, the hunt for black monkeys, several species of birds, and deers [18-22]. Exploitation activities done by local community is continuously carried out due to the absence of natural conservation efforts, it is feared to reduce the diversity of *O. croesus*, which will eventually lead to scarcity *O. croesus*. In the longer term can cause the extinction of *O. croesus*. The purpose of this research is to update the conservation status of *O. croesus* endemic butterflies in the Bacan Island based on the current information: the existence of the species, the characters and the habitat carrying capacity and conservation status.

## Experimental

This research was conducted in June until August 2015 at Sibela mountain natural conservation of Bacan Island at four locations. The location 1 is settlement, the location 2 is plantations, the location 3 is forest production, and the location 4 is isolated forest conversion.

### Materials

Tools and materials used are: GPS, camera, insect nets (sweep net), tweezers, altimeter, butterflies envelope paper.

### Methods

Methods of data collection in the field by making four transects (line-transect method) at each study site with 3 plot observations. Sampling was purposive random sampling. Data analysis was performed with:

a. Analysis of the existence of the butterfly using density analysis of individual species of butterflies, simple method of estimating density [23].

$$D = \ln n/n^2 \times n/m (n \cdot r^2) \times 10.000,$$

where is: *D* = density per hectare; *n* = total number of one individu; *n*<sup>2</sup> = total individu of species outside the radius of 25m; *m* = total observed points; *r* = radius (m).

b. Analysis of the characters and the carrying capacity of the habitat of butterflies by using a percentage (%) availability of food in each plot observation.

c. Analysis of butterfly conservation status is determined based on the standard of Regulation of the Minister of Environment and Forestry of the Republic of Indonesia [24] and the IUCN (*International Union for Conservation of Nature and Natural Resources*) the IUCN Red List on Wallace's Golden Birdwing Butterfly and *Ornithoptera croesus* in 2018 [25, 26] and the CITES (*Convention on International Trade in Endangered Species of Wild Fauna and Flora*) [27].

## Results and discussion

### *The existence of O. croesus endemic butterflies in Bacan Island*

*O. croesus* is an endemic butterfly in the Bacan Island of South Halmahera district that has aesthetic, economic, and high conservation values. The morphological descriptions *O. croesus* (Fig. 1).



Fig. 1. Species *O. croesus* males and females on Sibela mountain nature reserve area in Bacan Island




Fig. 2. Morphological structure of male and female *O. croesus* Endemic Butterfly of Bacan Island

To determine the existence of butterflies *O. croesus* on, in Sibela mountain natural conservation density data parameter butterfly species *O. croesus* were used on transect applied on the four observation sites with different heights (Table 1).


**Table 1.** Density of butterfly species at different elevations *O. croesus*

No	Location (m asl)	Density (individu/hectare)
1	20	20
2	200	14
3	400	11
4	800	14

Density of *O. croesus* at different elevation in Sibela mountain natural conservation has low density and a few individuals *O. croesus* were found in small quantities.

#### **Characters and Habitat Carrying Capacity**

The Region of Sibela mountain natural reservation is one of the endemic regions for the distribution of species of flora and fauna. In this area the *O. croesus* species live at a particular location on one of its habitat, especially at the edges of rivers or swamp which plant as food are available. Location and contour maps of the stew butterfly *O. croesus* habitat at Sibela mountain natural conservation (Fig. 3).



**Fig. 3.** Distribution of butterfly *O. croesus* on showed specific sidefor habitat/hotspot at Sibela mountain natural conservation of Bacan island.

The main hospot of *O. croesus* species on this region are located at a certain height of 20m above sea level (Lowland/settlement), 200m asl (Plantation Balitro), 400m (Forest production/River Ra), 800m asl (Forest finite conversion/Sibela talaga sago) or a buffer zone of Sibela mountain natural reservation, while at elevations above 800m asl butterfly *O. croesus* could not be found anymore. In general, a hotspot of *O. croesus* is marked by the availability of musaenda and asoka plantation as the food resource of *Ornithoptera* spp. Data presenting the percentage of food resource in each location (Table 2).

#### **Conservation Status *O. croesus* Endemic Butterflies in the Bacan Island**

Bacan Island is one of the spots in Wallace expedition. According to [28] Wallacea area included in the 25 "hotspots" most important for conservation. This area has 529 species of endemic invertebrates (1.9% of the total number in the world). Species are experiencing serious

## CONSERVATION STATUS OF THE ENDEMIC BUTTERFLY ORNITHOPTERA CROESUS

threat, because only 15% of the natural habitat that remains. Of the remaining natural habitat of these, 39.2% of which are in protected areas.

**Table 2.** Characteristics and habitat carrying *O. croesus* at the 4 locations

Location	Position(Coordinat)	Food plant Dominant	Cover (%)
Settlement (20m asl)	S = 00°37'21,8''	Musaenda	81
	E = 127°30'33,0''	Asoka	83
Plantation (200m asl)	S = 00°39'46,4''	Musaenda	78
	E = 127°31'46,1''	Asoka	71
Production forest (400m asl)	S = 00°40'08,6'' E = 127°32'29,9''	Musaenda	68
		Asoka	57
		Gusale	77
Conservation forest (800m asl)	S = 00°43'06,4'' E = 127°33'51,6''	Musaenda	53
		Gusale	81

Conservation status *O. croesus* at Sibela mountain natural conservation based on Red list Data Book of the IUCN (*International Union for Conservation of Nature and Natural Resources*) and Appendix II (Table 3) of CITES (*Convention on International Trade in Endangered Species of Wild Fauna and Flora*) [25, 26, 27].

**Table 3.** Conservation status of the endemic butterflies *O. croesus* in Bacan Island

Spesies Name	Government regulation (PP no 7 Th 1999 ) Conversion Status	IUCN Data Red Book Status	CITES Conversion Status	Endemic
<i>Ornithoptera croesus</i>	Protected	Vulnerable	Threat of extinction	√

The conservation status of endemic butterflies *O. croesus* in Bacan Island is protected and endangered. *O. croesus* is a kind of large winged butterfly most sought after by collectors because of the beauty of the color of its wings and is now listed as endangered and entered in Appendix II of CITES, thus the need for efforts to restrict international trade in captive specimens is a necessity. According to the Regulation of the Minister of Environment and Forestry of the Republic of Indonesia [24] it is known that *O. croesus* is a type of *O. croesus* butterfly that is protected because the population continues to decline. At this time, this butterfly became the object of hunting by collectors because of its wonderful shape and color patterns. In its natural habitat in the wild (out of conservation habitat) the survival of these butterflies are threatened by habitat loss as a place of life and reproduction, and as well as the shortage of feed due to conversion of forest lands.

Based on the analysis of the *O. croesus* conservation status above, it is necessary to build conservation of *O. croesus* butterflies to avoid extinction. Some conservation models that can be recommended for *O. croesus* are: *ex-situ* conservation models with captivity outside of their natural habitat. It is to accelerate the breeding of butterflies. These models consider the condition of the transfer functions of forests as *O. croesus* natural habitat is diminishing and positively correlated with the carrying capacity of the feed, so it can be assumed that with little amount of food available has caused the butterflies to migrate to obtain food. In addition, the *in-situ* conservation models also can be implemented by increasing the sustainability of the habitat through reforestation program at mount Sibela natural reservation and the reduction program over the function of forests into plantations and settlement.

The population of the endemic butterfly *O. croesus* in mount Sibela natural conservation has relatively low life span, it can be seen at each study site only a few individuals *O. croesus*

were found in relatively small quantities. In general, the diversity and abundance of butterfly populations in nature are influenced by the quality of the environment of its habitat. In areas with more natural diversity and abundance of butterflies higher than that of the areas that have been converted into agricultural land and plantations. The results of the study [29] showed that the diversity of the butterfly family Papilionidae at various altitudes on mount Sibela natural conservation of Bacan Island in all habitats (a height of 20 meters above sea level, 200 meters above sea level, 400 meters above sea level and 800 meters above sea level) were categorized as low as the value  $H'$  in the range of 1.5 to 3.5. *M. Indrawan, et al.*, [30] stated that generally the main threat to the population decline is due to the destruction of endemic species and habitat fragmentation, pollution, excessive utilization types, introduction of exotic species and the spread of disease. Furthermore [31-34] stated that the existence of butterflies in nature depend on the diversity of host plant, it is closely related to vegetation diversity in its habitat. Based on description above it can be said that the abundance/density of butterfly *O. croesus* is relatively low at mount Sibela natural conservation. This is likely to be influenced by the availability of forage plants other than environmental factors (temperature, humidity and light intensity), while the minimal availability of forage plants affected by conversion of forest lands.

The character and habitat capacity of the availability of forage plants for *O. croesus* at Sibela mountain natural reservation are categorized as quite abundant, so it can be said that Sibela mountain natural conservation of Bacan island owning sufficient food storage for the *O. croesus* butterfly. At each location forage plants: such as musaenda plant, asoka plant and gusale plant are found. *A. Shalihah, et al.*, [35] stated that the existence of the butterfly depends upon the availability of food resource both at lowland and highland habitat. The habitat has hostplant components and foodplant. Hostplant are the food resource for the larvae and the regular plants feed the adult butterfly.

Furthermore, [36, 37] suggests that the swallowtail butterflies species (*O. croesus*) live in the lowlands located in swamps and wet places (rivers). Butterflies generally are encountered at various altitudes, ranging from lowlands to highlands 0-2000 m above sea level and is more common found on tropical areas [38-41]. In this study, the distribution of *O. croesus* butterflies at Sibela mountain natural reservation is limited to a height of 800 m above sea level, it is influenced by physiological adaptation factors and the availability of forage plants for survival.

Conservation status of a species made over the years is based on the category of IUCN and CITES. Special IUCN make a list of all types of flora that needs special attention because of the potential in its natural habitat began to decline. IUCN also made a list of fauna, including animals that are traded internationally as rare and endemic species, while CITES has made a list of the types that need to be protected based on the degree of scarcity. This list is very helpful in efforts to prevent illegal logging, illegal wildlife trade and black market.

CITES divides species into categories based on the type of status scarcity in nature ie Appendix I of species already threatened with extinction so that the circulation between countries is prohibited, except for a specific purpose and does not damage the natural habitat. Appendix II includes species which have not endangered yet, but if international trade is not controlled then there is the risk of extinction. Whereas Appendix III contains species need to be supervised by a country internationally, even though the country to where the species type are available, has not yet require a mechanism tool to control internationally [42].

In Indonesia to anticipate the threat of natural resources damage, government issued various regulations containing procedures for setting in the utilization of natural resources in a way that still maintain the ecological balance. Some of the rules include: Regulation of the Minister of Environment and Forestry of the Republic of Indonesia [24] about types of plants and animals protected; [43] about preserving plants and animals; [44] about the use of wild plants and animals; [45] about conservation of living natural resources and their ecosystems and

[46], about utilization of types of plants and wildlife. About 25 species of butterfly that are protected by law mostly categorised as rare flora. The law also includes a conservation program *in-situ* and *ex-situ* specified to the endangered and rare type.

### Conclusions

The conservation status of *O. croesus* butterfly according to PP number 7 of 1999 is a protected animal, whereas according to IUCN the conservation status of *O. croesus* is vulnerable and according to CITES the conservation status of *O. croesus* is endangered. This is in accordance with the phenomenon in nature (natural habitat) of *O. croesus* butterflies in the Sibela mountain nature reserve with the condition of forest conversion causing habitat fragmentation which causes *O. croesus* butterflies as endemic butterfly species with low species density with sufficient environmental carrying capacity (availability of feed). The conservation efforts need to be made to the survival of *O. croesus* so that it is not extinct. Some conservation model recommendations for *O. croesus* are *ex situ* conservation by breeding around natural habitats and *in situ* conservation models through reforestation activities and reduction of forest conversion programs in the Sibela mountain reserve to increase environmental carrying capacity for the survival of *O. croesus* as endemic butterfly.

### Acknowledgments

The researchers would like to thank Mrs. Djunijanty Peggy, M.Sc. PhD from LIPI as butterfly researcher who has provided literature information, and Mukhlis, S.Pd for the support and assistance in the collection and identification of the *O. croesus* samples found in the conservation area of Sibela Mountain in Bacan Island.

### References

- [1] A. Mas'ud, *Diversity of Intraspecies of Ornithoptera croesus Endemic Butterflies of Bacan Island at Various Altitudes on Mount Sibela Based on Morphological Characters, Molecular Markings-RAPD and its Conservation Strategies and Development of Reference Books*, PhD Thesis of the UM Postgraduate Program, 2018, pp. 73-81.
- [2] A. Mas'ud, *Analysis of the level of knowledge of the communities around the area of the Sibela Mountain Nature Reserve against the Endemic Butterfly of Bacan Island (Study of Conservation Policy Material Ornithoptera Croesus)*, **Proceedings of the National Biology/Science Seminar and Learning, Malang State University**. ISBN; 978-602-73915-4-3, 2015, pp. 219-226.
- [3] L.P. Koh, N.S. Sodhi, B.W. Brook, *Ecological correlates of extinction proneness in tropical butterflies*, **Journal Conservation Biology**, **18**, 2004. pp. 1571-1578.
- [4] M.T. Irwin, P.C. Wright, C. Birkinshaw, B.L. Fisher, C.J. Gardner, J. Glos, M.J. Raherilalao, *Patterns of species change in anthropogenically disturbed forests of Madagascar*, **Biological Conservation**, **143**(10), 2010, pp. 2351-2362.
- [5] S.J. Willott, D.C. Lim, S.G. Compton, S.L. Sutton, *Effects of selective logging on the butterflies of a bornean rainforest*, **Conservation Biology**, **14**, 2000, pp. 055-1065.
- [6] J.K. Hill, K.C. Hamer, J. Tangah, M. Dawood, *Ecology of tropical butterflies in rainforest gaps*, **Journal Oecologia**, **128**, 2001, pp. 294-302.
- [7] K. Henle, K.F. Davies, M. Kleyer, C. Margules, Settele, *Predictors of species sensitivity to fragmentation*. **Biodiversity and Conservation**, **13**, 2004, pp. 207-251.
- [8] K.S. Bobo, M. Waltert, H. Fermon, J. Njokagbor, M. Muhlenberg, *Form forest to farmland butterfly diversity and habitat associations along a gradient of forest conversion in southwest cameroon*, **Jornal of Insect Consevation**, **10**, 2006, pp. 29-42.



- [9] W. Thuiller, G.F. Midgley, G.O. Hughes, B. Bomhard, G. Drew, M.C. Rutherford, F.I. Woodward, *Endemic species and ecosystem sensitivity to climate change in Namibia*, **Global Change Biology**, **12**(5), 2006, pp. 759-776.
- [10] D.F. Cleary, A.Ø. Mooers, *Burning and logging differentially affect endemic vs. widely distributed butterfly species in Borneo*, **Diversity and Distributions**, **12**(4), 2006, pp. 409-416.
- [11] S. Netherer, A. Schopf, *Potential effects of climate change on insect herbivores in European forests general aspects and the pine processionary moth as specific example*, **Forest Ecology and Management**, **259**(4), 2010, pp. 831-838.
- [12] I. Widhiono, *Diversity and abundance of java endemics butterfly (lepidoptera: rhopalocera) at Slamet Mountain, Central Java*, **Biospecies**, **7**, 2014, pp. 59-67.
- [13] G. Talavera, V. Dinca, R. Vila, *Factors affecting species delimitations with the GMYC model: insights from a butterfly survey*, **Methods in Ecology and Evolution**, **4**(12), 2013, pp. 1101-1110.
- [14] M. Pacifici, W.B. Foden, P. Visconti, J.E. Watson, S.H. Butchart, K.M., Kovacs, R.T. Corlett, *Assessing species vulnerability to climate change*, **Nature Climate Change**, **5**(3), 2015, pp. 215-224.
- [15] F.V. Florens, J.R. Mauremootoo, S.V. Fowler, L. Winder, C. Baider, *Recovery of indigenous butterfly community following control of invasive alien plants in a tropical island's wet forests*, **Biodiversity and Conservation**, **19**(14), 2010, pp. 3835-3848.
- [16] A. Salisbury, J. Armitage, H. Bostock, J. Perry, M. Tatchell, K. Thompson, Editor's Choice: *Enhancing gardens as habitats for flower-visiting aerial insects (pollinators): should we plant native or exotic species?*, **Journal of Applied Ecology**, **52**(5), 2015, pp. 1156-1164.
- [17] T. Olivier, R. Schmucki, B. Fontaine, A. Villemey, F. Archaux, *Butterfly assemblages in residential gardens are driven by species' habitat preference and mobility*, **Landscape Ecology**, **31**(4), 2016, pp. 865-876.
- [18] A. Mas'ud, Sundari, *Study of Soil Epifauna Community Structure in the Mount Sibela Halmahera Selatan Conservation Area in North Maluku*, **Bioedukasi**, **2**(1), 2011, pp. 1-9.
- [19] M.A.R. de Mattos Vieira, E.M. von Muhlen, Jr, G.H. Shepard, *Participatory monitoring and management of subsistence hunting in the Piagaçu-Purus reserve, Brazil*, **Conservation and Society**, **13**(3), 2015, pp. 254-264.
- [20] L. Parry, C.A. Peres, *Evaluating the use of local ecological knowledge to monitor hunted tropical-forest wildlife over large spatial scales*, **Ecology and Society**, **20**(3), 2015, Article Number: 15.
- [21] K.M. Fleisher, *The distribution, habitat use, and conservation status of three Atlantic Forest monkeys (*Sapajus xanthosternos*, *Callicebus melanochir*, *Callithrix sp.*) in an agroforestry/forest mosaic in Southern Bahia, Brazil*, **International Journal of Primatology**, **36**(6), 2015, pp. 1172-1197.
- [22] R. Sreekar, G. Huang, J.B. Zhao, B.O. Pasion, M. Yasuda, K. Zhang, R.T. Corlett, *The use of species-area relationships to partition the effects of hunting and deforestation on bird extirpations in a fragmented landscape*, **Diversity and Distributions**, **21**(4), 2015, pp. 441-450.
- [23] C. Bibby, N.D. Burges, D.A. Hill, **Bird Census Techniques**, University Press, Cambridge, 1992, p. 91.
- [24] \* \* \*, Regulation of the Minister of Environment and Forestry of the Republic of Indonesia Number P.106/MENLHK/SETJEN/KUM.1/12/2018 Regarding the Second Amendment to the Regulation of the Minister of Environment and Forestry Number P.20/MENLHK/ SETJEN/KUM.1/6/2018 About the Types of Plants and Animals Protected, Minister of Environment and Forestry of the Republic, 2018, p. 26-27.

- [25] \* \* \*, **The International Union for Conservation of Nature Red List of Threatened Species (Wallace's Golden Birdwing Butterflies)**, IUCN Redlist, 2018. <https://www.iucnredlist.org/species/15517/727365>. Visited on 25 April, 2020.
- [26] M. Bohm, *Ornithoptera croesus*, **The IUCN Red List of Threatened Species 2018**: e.T15517A727365. <https://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T15517A727365.en>, 2018. Downloaded on 25 April 2020.
- [27] \* \* \*, **Convention on International Trade in Endangered Species of Wild Fauna and Flora**, CITES, p. 52. 2019. <https://www.cites.org/eng/app/appendices.php>. Visited on 25 April 2020.
- [28] N. Myers, R.A. Mittermeier, C.G. Mittermeier, G.A. Da Fonseca, J. Kent, *Biodiversity hotspots for conservation priorities*, **Nature**, **403**, 2000, pp. 853-858.
- [29] A. Mas'ud, S. Hasan, A. Abdullah, *The family Papilionidae Butterfly Diversity in Different Altitude Region Sibela Mountains Nature Reserve*, **Proceedings of the National Seminar VI Biodiveritas Department of Biology**, **3**, 2016, pp. 237-243.
- [30] M. Indrawan, R.B. Primack, J. Supriatna, **Conservation Biology**, revised edition. Obor Indonesia, Jakarta, 2007, p. 89.
- [31] K.C. Hamer, J.K. Hill, S. Benedick, N. Mustaffa, T.N. Sherratt, M.T. Maryati, *Ecology of butterflies in natural and selectively logged forests of northern Borneo: The importance of habitat heterogeneity*, **Journal of Applied Ecology**, **40**(1), 2003, pp. 150-162.
- [32] J. Krauss, I. Steffan-Dewenter, T. Tschardt, *Landscape occupancy and local population size depends on host plant distribution in the butterfly *Cupido minimus**, **Biological Conservation**, **120**(3), 2004, pp. 355-361.
- [33] R.L. Dennis, *Butterfly habitats, broad-scale biotope affiliations, and structural exploitation of vegetation at finer scales: the matrix revisited*, **Ecological Entomology**, **29**(6), 2004, pp. 744-752.
- [34] M. Syafitri, D.R. Untari, J.I.M. Sari, I. Arifin, **Diversity and distribution of type butterflies (Lepidoptera) di resort mountain putri, national park of mount Gede Pangrango**, Departement of Biology, State University of Jakarta, 2010, p. 75.
- [35] A. Shalihah, Pamulang, R. Cindy, W. Riskawati, Z. I. Anwar, **Butterfly Jatnangor Padjadjaran University Campus**. HMDP UNPAD, 2012, p. 89.
- [36] E.V. Herkenhoff, R.F. Monteiro, A.P. Esperanco, A.V.L. Freitas, *Population biology of the endangered fluminense swallowtail butterfly *Parides ascanius* (Papilionidae: Papilioninae: Troidini)*, **The Journal of the Lepidopterists' Society**, **67**(1), 2013, pp. 29-34.
- [37] S. Safian, *Observation of hill-topping behaviour by the Giant African Swallowtail-*Papilio antimachus* Drury, 1782 and other recent records from Liberia (West Africa)(Lepidoptera: Papilionidae)*, **SHILAP Revista de Lepidopterologia**, **41**(163), 2013, pp. 323-329.
- [38] T.W. Pyrcz, J. Wojtusiak, *The vertical distribution of pronophiline butterflies (Nymphalidae, Satyrinae) along an elevational transect in Monte Zerpa (Cordillera de Mérida, Venezuela) with remarks on their diversity and parapatric distribution*, **Global Ecology and Biogeography**, **11**(3), 2002, pp. 211-221.
- [39] I. Widhiono, *Diversity of butterflies in four different forest types in Mount Slamet, Central Java, Indonesia*, **Biodiversitas Journal of Biological Diversity**, **16**(2), 2015, pp 15-21.
- [40] L.V. Vu, T.C. Bonebrake, M.Q. Vu, N.T. Nguyen, *Butterfly diversity and habitat variation in a disturbed forest in northern Vietnam*, **The Pan-Pacific Entomologist**, **91**(1), 2015, pp. 29-38.
- [41] B. K. Acharya, L. Vijayan, *Butterfly diversity along the elevation gradient of eastern Himalaya, India*, **Ecological Research**, **30**(5), 2015, pp. 909-919.
- [42] T. Soehartono, A. Mardiastuti, **CITES Implementation in Indonesia**. Nagao Natural Environment Foundation, 2002, p. 89.

- [43] \* \* \*, **Republic of Indonesia Government Regulation (PP) Number 7 of 1999 (7/1999) about Preserving Plants and Animals**, President of the Republic of Indonesia, State Minister of the Republic of Indonesia, 1999, pp. 1-24.
- [44] \* \* \*, **Republic of Indonesia Government Regulation (PP) No. 8 of 1999 Concerning the Use of Wild Plants and Animals**, President of the Republic of Indonesia, State Minister of the Republic of Indonesia, 1999, pp. 1-25.
- [45] \* \* \*, President of the Republic of Indonesia. Law of the Republic of Indonesia (UURI) No. 5 of 1990 concerning **conservation of living natural resources and their ecosystems**. State Minister of the Republic of Indonesia, 1990, pp. 1-12
- [46] \* \* \*, Government Regulation of the Republic of Indonesia Number 8 of 1999 concerning **utilization of types of plants and wildlife**, President of the Republic of Indonesia, State Minister of State of the Republic of Indonesia, 1999, pp. 1-20.
- 

*Received: December 16, 2018*

*Accepted: August 24, 2020*

# Conservation Status of Ornithoptera Croesus, Endemic Butterfly of Bacan Island

## ORIGINALITY REPORT

5%

SIMILARITY INDEX

3%

INTERNET SOURCES

1%

PUBLICATIONS

7%

STUDENT PAPERS

## PRIMARY SOURCES

1

[repository.uin-malang.ac.id](https://repository.uin-malang.ac.id)

Internet Source

2%

2

Submitted to Universitas Brawijaya

Student Paper

1%

3

Submitted to Universitas Diponegoro

Student Paper

1%

4

Abdu Mas`ud, Sundari ., Nurhasanah ., Chumidach Roini, Mohamad Amin, Fatchur Rohman, Alisi .. "Genetic Variability of Ornithoptera croesus toeantei Endemic Butterfly in Morotai Island, Based on Morphology and Molecular", Journal of Entomology, 2019

Publication

1%

Exclude quotes

Off

Exclude matches

< 1%

Exclude bibliography

On